

Therapiestudien für akute Leukämien und MDS Stand 07.2020

Primärtherapie

ALL / LBL				APL	AML (non-APL)						MDS
GMALL-Register Spiekermann, Sand, Reischer, Rückel, Blumenberg, Rejeski				AMLCG-Register [Spiekermann, Grunwald, Bücklein, Reischer, Blumenberg]							
18 - 55J		>55J		medically fit			medically unfit				
BCR/ABL neg.	BCR/ABL pos.	BCR/ABL neg.	BCR/ABL pos.	APOLLO Spiekermann, Metzler, Plohmman <u>Indikation:</u> Hochrisiko-APL <u>Design:</u> ATRA+ATO vs. ATRA+Chemo	MOSAIC Subklewe, Bücklein <u>Indikation:</u> 18-75J CBF+ und/oder FLT3- TKD+/-ITD+ AML <u>Design:</u> Phase I/II 7+3 ± Mido ± GO In Planung	BLAST Subklewe, Reischer <u>Indikation:</u> 18-75J (nur <60J: non-favorable), AML in CR/CRi/CRp nach Induktionstherapie <u>Design:</u> Ph III Kons. HiDAC +/- BL-8040 Rekrutierung pausiert	ETAL-1 Tischer <u>Indikation:</u> Pat. (≤60 J) in CR1 mit int. Risiko & HLA-komp. Spender. CAVE: Frühzeitiger Einschluss! <u>Design:</u> alloTx vs. Chemotherapie	AG 120 Subklewe, Metzler <u>Indikation:</u> AML IDH1 mut. <u>Design:</u> Azacitidine +/- AG-120	AML1003 Subklewe, Bücklein <u>Indikation:</u> untreated AML <u>Design:</u> Phase I: Cusatumumab+Aza/ +Ven/+Ven/Aza	AARON Subklewe, Bücklein <u>Indikation:</u> AML > 65J <u>Design:</u> Phase I/II Aza + Nivo + Relatlimab In Planung	
GMALL 08/2013 Spiekermann, Subklewe, Reischer, Sand, Edlinger, Banck, Rückel, Cusan <u>Indikation:</u> ALL/LBL <u>Design:</u> Therapieoptimierung											
EWALL-BOLD Subklewe, Bücklein <u>Indikation:</u> ALL/LBL <u>Design:</u> Chemo + Blina											
GMALL-MOLACT1-BLINA Subklewe, Bücklein <u>Indikation:</u> CD19+ B-ALL in CR, mol. MRD >10 ⁴ <u>Design:</u> Einarmig Blinatumomab											